

APPLICATION UNDER UNITED STATES PATENT LAWS

Atty. Dkt. No. PM 265159/PHM70237/UST
(M#)

Invention: USE OF 2'-O-METHYL RNA AS HYBRIDISATION PROBE

Inventor (s): THEAKER, Jane
CALLAGHAN, Kay

Pillsbury Madison & Sutro LLP
Intellectual Property Group

Attorneys
Telephone:

This is a:

- ☐ Provisional Application
- ☐ Regular Utility Application
- ☐ Continuing Application
- ☒ PCT National Phase Application
- ☐ Design Application
- ☐ Reissue Application
- ☐ Plant Application
- ☐ Substitute Specification

Sub. Spec Filed
in App. No. _____ / _____

SPECIFICATION

USE OF 2'-O-METHYL RNA AS HYBRIDISATION PROBE

This invention relates to improvements of the nucleic acid probes known as Molecular Beacons. Specifically, the use of modified nucleic acids for the synthesis of such probes has advantages in the detection of nucleic acid targets, in several applications including in situ hybridisation and homogeneous real time PCR assays.

Molecular Beacons are recently described nucleic acid probes that fluoresce upon hybridisation (Tyagi and Kramer, Nature Biotechnology 14: 303-308; 1996). The secondary structure of the probes is key and depends upon their sequence. Each probe has a target complementary portion (the loop) of about 15 or more nucleotides and additional sequences appended to each end that are complementary to each other. Finally, the probe has a fluorescent reporter molecule on one end and a fluorescence quencher at the other end. The self complementary end sequences can form stems by base pairing, thus bringing the fluorophore and quencher into close proximity and eliminating the fluorescence signal. However, in the presence of (loop) complementary target, the loop and target base pair and this "stretches out" the probe strand, thus removing the fluorophore from the quencher and allowing the reporter to fluoresce when illuminated appropriately. In this way, fluorescence is dependent upon hybridisation.

Molecular Beacons offers a number of advantages. It can be performed as a homogeneous assay. Because the "switching on" of the signal only occurs in the presence of target, there is no requirement to wash away excess unbound probe. This makes Molecular Beacons ideal for applications such as *in situ* hybridisation and real time PCR. It yields low background signals. In the absence of target and at an appropriate temperature, the probe self anneals and quenches fluorescence. When amplicon accumulates, some of the probe binds and is unwound, thereby generating fluorescence. In this way homogeneous (no wash, closed tube) fluorescence is produced and the low signals in the absence of target results in high ratios of signal to noise. It offers design flexibility. The use of standard stems allows rapid and reliable probe design. Furthermore, by correct design and optimisation of the stems, backgrounds at temperatures appropriate for PCR can be minimised.

For homogeneous analysis, the loop element of a Molecular Beacon should be thermodynamically favoured over the stem portion, but the stem should form readily at the

assay temperature. These key requirements make it likely that long loops and stems are necessary, which has consequences for the synthesis yields and purity of Molecular Beacons. Furthermore, while long loops produce good separations between fluorophore and quencher, long stems melt over a broad temperature window leading to higher background signals and greater noise in those signals.

We have devised and now provide modified RNA probes which have a number of beneficial characteristics when used as Molecular Beacons.

Therefore in a first aspect of the present invention we provide a Molecular Beacons assay method using a 2'-O-substituted RNA probe having both a donor and a quencher species attached. The 2'-O-substituent is conveniently a methyl, propyl, butyl or allyl group. It is preferably a 2-methyl group.

The Molecular Beacons assay method is disclosed in WO-95/13399 (Public Health Research Institute of New York). This essentially comprises the detection of a pre-selected nucleic acid sequence by contacting a sample believed to contain said sequence with a Beacons probe under hybridisation conditions such that there is a detectable change in Beacons signal from the probe if the pre-selected sequence is present in the sample.

The Beacons probe is a unitary probe comprising a single stranded target complementary sequence, a stem duplex consisting of nucleotide sequences 5' and 3' to the target complement sequence and having a melting temperature lower than the target complementary sequence/target sequence melting temperature, and at least one label pair, each pair comprising a first label conjugated to the probe at or near the 5' terminus of the probe and a second label conjugated to the probe at or near the 3' terminus of the probe. Under assay conditions, hybridisation of the target complementary sequence to the target sequence leads to a change in Beacons signal from the label pair.

The donor and acceptor species are attached to the Molecular Beacons probe in any convenient way. By way of non-limiting example either species may be attached to the 3' terminus of the probe via controlled pore glass (CPG) based synthesis or attached to the 5' terminus via 5'-phosphoramidite chemistry. The donor and quencher species are attached at any convenient locations on the probes, such as for example at or near the ends of the probe, preferably at the ends of the probe. Examples of convenient donor species will be apparent to the scientist of ordinary skill and include FAM, TET, JOE, HEX, ROX, BODIPY and EDANS. Convenient acceptor species include TAMRA, Pyrene butyrate, and DABCYL.

Still further convenient details are found in, for example Livak et al. "PCR methods and applications, 1995, 4, 357-362; WO-95/13399 (Public Health Research Institute of New York).

Advantages of the 2'-O-substituted probes include a higher T_m when annealed to DNA target, relative to the "standard" DNA probes; this means that shorter probes can be designed to function at similar temperatures and therefore a higher level of specificity is possible - a single mismatch within a short oligonucleotide has a greater destabilising effect than when the same mismatch is in a larger hybridised region

Secondly 'traditional' Molecular Beacons perform optionally in magnesium concentrations of 3.5 mM or greater. With the new 2'-O-substituted Molecular Beacons significantly lower magnesium concentrations can be utilised. This has substantial benefits for the specificity of PCR and in particular ARMS reactions. [ARMS technology is described in, for example Newton et al., 1989, Nucleic Acids Research (17) 2503-2516].

Also, annealing of 2'-O-substituted RNA to complementary 2'-O-substituted RNA targets is even more favourable than to DNA targets; this permits a shorter stem region to be used which has particular benefits for the design of effective Beacon probes. This has three functional benefits for the use of Molecular Beacons, particularly in real time PCR systems.

Firstly, when a Molecular Beacon is cooled from high temperature (94°C) to low temperature (20°C) the fluorescence changes from high (Beacon is a random coil) to low (Beacon has adopted a stem/loop formation), see Figure 1. The temperature range of the transition from "random coil" to stem-loop structure depends upon the length of the stem sequence; longer sequences are expected to melt over a wider temperature window, possibly in a series of small steps, while shorter stems are more likely to undergo a single, rapid stem transition. This is important for the design of effective real time assays in which backgrounds should be as low as possible: the stem/loop structure should be fully formed at the temperature at which the fluorescence is measured and the probe/target interaction should also be favoured at this temperature.

Secondly, short stems have an additional advantage in that they should exhibit lower noise in positive hybridisations, due to their lower flexibility, and hence reduced ability to give a degree of fluorophore quenching.

Thirdly, at lower temperatures, the free stem sequences can "find" each other, even when the loop and the target have annealed. The hybridised duplex can be bent around and

thus quenching of the fluorescence signal can occur. The consequences of this feature also favour the design of Molecular Beacons with shorter loop portions to minimise this hybrid bending, since shorter duplexes are more physically constrained and cannot bend so easily into this closed formation

5 A still further advantage is that where true quantitation of PCR product is required, it is desirable to have no cleavage of dual labelled probes since the fluorescence observed in such reactions reflects the accumulation of the cleavage reactions through all the previous cycles, thus obscuring to some extent the true level of amplicon actually accumulated thus far. Many of the enzymes commonly used in PCR have an endogenous 5'-nuclease activity and
10 may cleave the probes

Therefore in a further aspect of the present invention we provide 2'-O-substituted RNA probes having both a donor and a quencher species attached. The 2'-O-substituent is conveniently a methyl, propyl, butyl or allyl group. It is preferably a 2-O-methyl group. These modified nucleic acids are nuclease resistant; this is important in assays where nucleases may
15 be present such as in situ hybridisation or in real time PCR assays.

The invention further relates to diagnostic kits comprising one or more of the Beacons probes of the invention, together with appropriate buffers and other reagents, and instructions for use.

The invention will now be illustrated but not limited by reference to the following
20 Example and Figures wherein:

Figure 1 shows cooling curves for 2-Methyl RNA Molecular Beacon 0007M, in the presence or absence of synthetic oligonucleotide target (R297). Fluorescence readings were taken throughout the cooling range and plotted against temperature. Raw fluorescence
25 intensity is along the Y-axis.

Figure 2 shows a similar series of curves in which PCR amplicons at various concentrations from neat to 1/8 diluted were cooled in the presence of Beacon 0007M. On the X-axis temperature decreases from 94°C to 11°C as before, but to permit accurate tube to tube comparisons, the data are equalised to a baseline between temperatures 84°C to 74°C where
30 each of the curves is flat. When the mixtures have cooled to below 50°C, the curves reach their minimum fluorescences and it is appropriate to compare them. The fluorescence value at this point rise with the quantity of target within the tube.

Figure 3 illustrates the increase in Beacon fluorescence monitored at the same point in every PCR cycle (the 60°C anneal step). Where template is included in the reaction, fluorescence increases above the background (shown by the no template control). This increase becomes significant at ~28 cycles.

Figure 4 shows the same reactions using Beacon 0009M in place of the previous 0007M Beacon. The fluorescence grows smoothly with less background noise and reaches substantially larger values. This reflects the improved efficiency of this Beacon due to enhanced design.

Figure 5 shows the detection of normal alleles (N mix) of the hereditary haemochromatosis mutations C282Y het and C282Y homo, together with wild type and control sequences

Figure 6 shows the detection of mutant alleles (M mix) of the hereditary haemochromatosis mutations C282Y het and C282Y homo, together with wild type and control sequences

Figure 7 shows the detection of normal alleles (N mix) of the hereditary haemochromatosis mutation H63D, together with wild type and control sequences.

Figure 8 shows the detection of mutant alleles (M mix) of the hereditary haemochromatosis mutation H63D, together with wild type and control sequences.

Example 1

Reagents:

Primers and Probes

R351: CGC TGA TGA ATG TGA AAA ATC TAA

R352: AGA AGT TCC AGA TAT TGC CTG CTT

0007M (2-methylRNA Molecular Beacon): (FAM)-GCG AGC AAA AGA CCU AUU AGA

CAC AGA GAA GCU CGC-(Quencher); underlined portions indicate the self complementary stems.

R297: CTT TTG TTC TCT GTG TCT AAT AGG TCT TTT TCT GAA; synthetic target for Beacon 0007M. underlined region is the complementary portion.

Other Reagents:

10xARMS(35) Buffer: 100 mM Tris-HCl (pH 8.3 at 25°C), 500 mM KCl,
3.5 mM MgCl₂, 0.1% gelatin)

1 mM dNTPs: 1mM each dNTP diluted from
5 10mM stocks(Pharmacia)

Amplitaq Gold (5U/μl): from Perkin-Elmer

ROX standard: ROX conjugated oligonucleotide, 600 nM stock solution

Melt Characteristics:**Reaction Mix**

10 400 nM Molecular Beacon 0007M in 1xARMS (3.5) [diluted from 10-fold stock (10 x ARMS)] plus dNTPs at 100mM final, ROX standard at 60 nM and targets at various concentrations. Targets are either synthetic oligo target (R297) at 1μM or double stranded amplicon produced by PCR with primers R351 and R352 and serially diluted.

Cycling Parameters

15 94°C 2 min; then 84 steps of 15 seconds decreasing from 94°C to 11°C using an ABI PRISM system 7700, fluorescence readings monitored at each temperature.

Results

Figure 1 shows the results obtained with 0007M in the presence (◆) or absence (■) of target. The high fluorescences at elevated temperatures show the stem has not formed and the probe is essentially randomly coiled. As the temperature decreases, the fluorescence decreases in the "no target" reaction, while in the presence of excess target, an increase in fluorescence is observed, peaking at around 56°C. The subsequent reduction in this fluorescence reflects the high affinity of the stems for each other and at lower temperatures, the probe target duplex can be bent permitting stems to form and fluorescence to switch off.

25 Figure 2 shows a similar series of curves with varying quantities of double-stranded amplicon: a 2-fold serial dilution from "neat" to 1/16 and a negative control. The data have been equalised for fluorescence between 88°C and 74°C (where each line is flat) to allow direct comparison between tubes. There is a clear difference between positive and negative samples and the dilution series of target numbers is clearly reflected in the relative

fluorescences, particularly at lower temperatures. There are two further observations to be drawn from this data:

1. the larger amplicon target is less prone to subsequent bending and closure than the oligonucleotide target;
2. the target was heat denatured only once and it might have been expected that the "other" strand would displace the probe from its binding site. This was not the case since fluorescences remained substantial throughout the cooling.

Amplification Reactions:

Reaction Mix

1xARMS (3.5), buffer added to 100 μ M dNTPs, 500 nM each primer (R351, R352), 60 nM ROX standard, 400 nM Beacon 0007M, 2 Units Amplitaq Gold (per 50 μ l reaction), 5 μ l (50 ng) human genomic DNA.

Cycling Conditions

The reactions were cycled and fluorescence read using the conditions below.
94°C for 20 min to activate the Amplitaq Gold. Followed by 40 cycles of:
94°C for 41s, 60°C for 42 s, 72°C for 52 s. Fluorescence data was collected and analysed for the 60°C anneal step.

Results

The output from an amplification reaction as described above is shown in Figure 3.
There is a clear difference between positive and negative samples using this system. We anticipate that the probe design and reaction conditions may be further optimised, thus yielding increased fluorescences, whilst minimising the signal noise and enhancing the sensitivity of the technique.

Example 2

In this Example we disclose a further, improved Molecular Beacon. In this the loop portion of the probe was enlarged to ensure strong binding to target and good physical separation of fluorophore from quencher. In addition, the stem was decreased in size to a CGCG tetramer which has the multiple benefits of short length to minimise the folding back

- 8 -

of the bound duplex, high T_m due to its particular sequence composition, 5'-terminal C adjacent to the fluorophore which avoids possible quenching by the G, as present in probe 0007M.

The improved Beacon is 0009M: (FAM) - CGC GGA AAA ASA CCU AUU AGA
 5 CAC AGA GAA CAC GCG - (Quencher). Underlined portions are the stem. All bases are 2'-O-methyl RNA.

Example 3

10 **Detection of Hereditary Haemochromatosis mutations using 2'-O-methyl RNA molecular beacons**

1) C282Y ARMS Test

15 **Materials and Methods**

Two mixes were made up, the first (N) to detect the normal (wild type) allele and the second (M) to detect the mutant allele. The mixes differ only by the 3' base of the ARMS primer. Final concentrations of mix components are shown below:

20 C282Y N mix	C282Y M mix
1x Beacons buffer	1x Beacons buffer
100mM dNTPs	100mM dNTPs
1x Passive Reference (60nM ROX)	1x Passive Reference (60nM ROX)
0.5mM common forward primer (R279-97)	0.5mM common forward primer (R279-97)
0.5mM reverse ARMS normal primer (R280-97)	0.5mM reverse ARMS normal primer (R281-97)
0.4mM molecular beacon (0037M)	0.4mM molecular beacon (0037M)
1 unit AmpliTaq Gold enzyme	1 unit AmpliTaq Gold enzyme

Composition of 1x Beacons buffer:

50mM KCl

10mM Tris (pH 8.3)

- 9 -

3.5mM MgCl₂

0.01% (w/v) gelatin

Primer / Beacon Sequences (5' → 3'):

R279-97 AAGTGCCTCCTTTGGTGAAGCTGACACA

5 R280-97 TGATCCAGGCCTGGGTGCTCCACCTGAC

R281-97 TGATCCAGGCCTGGGTGCTCCACCTGAT

0037M (FAM)-CGCGAGUUCGAACCUAAAGACGUAUUGCCCAACGCG-
(Quencher)

10 The mixes were dispensed into 20µl aliquots in optical tubes. 5µl of genomic DNA (prepared from blood by alkali lysis and diluted 1/5 in water) was added to duplicate aliquots of both mixes. Examples of wild type, heterozygous mutant and homozygous mutant samples were used.

15 The tubes were placed in a thermal cycler (Applied Biosystems 7700) and the following PCR program was run:

20 minutes at 94°C followed by

20 cycles of

45 seconds at 94°C

45 seconds at 60°C

20 then 25 cycles of

45 seconds at 94°C

45 seconds at 40°C

The results are shown in Figure 5 and Figure 6

25 The results above demonstrate that the N mix only amplifies genomic DNA which contains the wild type allele (solid diamonds and clear triangles) and the M mix only amplifies genomic DNA which contains the mutant allele (clear triangles and solid circles). Amplification does not occur in either mix in the absence of genomic DNA (Xs).

2) H63D ARMS Test

30 Materials and Methods

Two mixes were made up, the first (N) to detect the normal (wild type) allele and the second (M) to detect the mutant allele. The mixes differ only by the 3' base of the ARMS

primer. All primers in the H63D test have the same 26mer non-homologous tail sequence at their 5'-end. Final concentrations of mix components are shown below:

H63D N mix	H63D M mix
1x Beacons buffer	1x Beacons buffer
100mM dNTPs	100mM dNTPs
1x Passive Reference (60nM ROX)	1x Passive Reference (60nM ROX)
0.5mM common forward primer (R369-98)	0.5mM common forward primer (R369-98)
0.5mM reverse ARMS normal primer (R370-99)	0.5mM reverse ARMS normal primer (R371-98)
0.4mM molecular beacon (MB026-98)	0.4mM molecular beacon (MB026-98)
1 unit AmpliTaq Gold enzyme	1 unit AmpliTaq Gold enzyme

5 Composition of 1x Beacons buffer

50mM KCl

10mM Tris (pH 8.3)

3.5mM MgCl₂

0.01% (w/v) gelatin

10 Primer / Beacon Sequences (5'→3')

R369-98

CGGTACTAGCGTACCACGTGTCGACTTCCTACTACACATGGTTAAGGCCTG

R370-98

CGGTACTAGCGTACCACGTGTCGACTGGGCTCCACACGGCGACTCTCAAG

15 **R371-98**

CGGTACTAGCGTACCACGTGTCGACTGGGCTCCACACGGCGACTCTCAAC

MB026-98 (FAM)-CGCGGGAUGACCAGCUGUUCGUGUUCUACGCG-(Quencher)

The mixes were dispensed into 20μl aliquots in optical tubes. 5μl of genomic DNA (prepared from blood using the Gentra PureGene kit and diluted in water to 10ng/μl) was added to duplicate aliquots of both mixes. Examples of wild type, heterozygous mutant and homozygous mutant samples were used.

- 11 -

The tubes were placed in a thermal cycler (Applied Biosystems 7700) and the following PCR program was run:

20 minutes at 94°C followed by

20 cycles of

45 seconds at 94°C

45 seconds at 60°C

then 25 cycles of

45 seconds at 94°C

45 seconds at 40°C

The results are shown in Figures 7 and 8.

Again, the results demonstrate that the N mix only amplifies genomic DNA containing the wild type allele (solid diamonds and clear triangles) and the M mix only amplifies genomic DNA containing the mutant allele (clear triangles and solid circles). Amplification does not occur in either mix in the absence of genomic DNA (Xs).

The amplification protocol was identical to the run shown in Figure 3. The results from this amplification are shown in Figure 4.

-12-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5 (i) APPLICANT:

(A) NAME: Zeneca Limited

(B) STREET: 15 Stanhope Gate

(C) CITY: London

(D) STATE: England

(E) COUNTRY: United Kingdom

(F) POSTAL CODE (ZIP): W1Y 6LN

(G) TELEPHONE: 0171 304 5000

(H) TELEFAX: 0171 304 5151

(I) TELEX: 0171 304 2042

15 (ii) TITLE OF INVENTION: ASSAYS

(iii) NUMBER OF SEQUENCES: 13

20 (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

25

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: GB 9715522.0

(B) FILING DATE: 24-JUL-1997

30

-13-

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CGCTGATGAA TGTGAAAAAT CTAA

24

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

AGAAGTTCCA GATATTGCCT GCTT

24

-14-

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GCGAGCAAAA GACCUAUUAG ACACAGAGAA GCUCGC

36

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CTTTTGTCT CTGTGTCTAA TAGGTCTTTT TCTGAA

36

-15-

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CGCGGAAAAA ACCAAGACAC AGAGAACACG CG

32

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

AAGTGCCTCC TTTGGTGAAG CTGACACA

28

-16-

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

TGATCCAGGC CTGGGTGCTC CACCTGAC

28

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

TGATCCAGGC CTGGGTGCTC CACCTGAT

28

-17-

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

CGCGAGUUCG AACCUAAAGA CGUAUUGCCC AACGCG

36

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 51 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

GCGTACTAGC GTACCACGTG TCGACTTCCT ACTACACATG GTTAAGGCCT G

51

-18-

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GCGTACTAGC GTACCACGTG TCGACTGGGC TCCACACGGC GACTCTCAAG

50

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

GCGTACTAGC GTACCACGTG TCGACTGGGC TCCACACGGC GACTCTCAAC

50

-19-

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

CGCGGGAUGA CCAGCUGUUC GUGUUCUACG CG

32

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167